
APPENDIX C. AIR QUALITY SYNTHESIS

Annie Esperanza, Sierra Nevada Network Air Quality Specialist, Sequoia and
Kings Canyon National Parks

Liz van Mantgem, Physical Sciences Technician, Sequoia and Kings Canyon
National Parks

September 2005

General Description	3
Criteria Pollutants: Description and Issues.....	7
Attainment.....	10
Pollutant.....	11
Federal	11
State	11
Air Quality in Sierra Nevada Parks.....	12
Air Resources Division of the National Park Service.....	24
Monitoring and Research Needs	31
References.....	36

APPENDIX C. AIR QUALITY SYNTHESIS

General Description

Air pollution is currently recognized as one of the most significant threats to natural resources of the Sierra Nevada. A description of these threats can be found, *supra*, in the main body of this Monitoring Plan (section 1.3.4.4).

Sources of air pollutants are local, regional, and in some instances, global. The National Park Service (NPS), in cooperation with state and other federal agencies, is making concerted efforts to reduce resource damage caused by air pollution. This is done by an intensive monitoring program, support for research, identifying and seizing opportunities to participate in decisions being made by regulatory groups, and expanding the education of park visitors and those concerned about the future of these Sierra Nevada parks.

Kings Canyon, Sequoia, and Yosemite National Parks are designated Class I air sheds under the Clean Air Act (1977 amendment). As such, the parks are afforded the greatest degree of air quality protection under the Clean Air Act, and the National Park Service is required to do all it can to ensure that air quality related values are not adversely affected by air pollutants. This includes participation in the review of permits of those sources whose emissions will potentially affect the parks. Devils Postpile National Monument, the fourth NPS unit in the Sierra Nevada Network (SIEN), is listed as a Class II air shed, which still has a mandate to protect air quality, but it is not as stringent as that required for parks with Class I air sheds.

Air pollution impairs visibility and injures plant and animal life. The once vast panoramas from vista points in the parks looking westward are now highly obscured by regional haze. Plant species differ in their sensitivity to pollutants; there are numerous species within SIEN parks that are damaged by air pollutants. For example, studies have shown that Jeffrey and ponderosa pines are especially sensitive to ozone (Duriscoe and Stolte 1989). Ozone-related effects include foliar damage, reduced needle retention, and reduced growth.

Giant sequoia seedlings show needle damage at current ozone levels. Acid deposition has been found to affect the chemical composition of lakes and streams within the parks, which can harm aquatic life. Contaminants used in the nearby San Joaquin Valley are found in rain and snow samples within the parks, raising concern over potential impacts to both aquatic and terrestrial wildlife.

The San Joaquin Valley, west of the Sierra Nevada parks, is a trap for air pollutants originating in the Valley as well as from pollutants from cities along the central California coast. These pollutants are carried in prevailing winds. Southward wind patterns carry pollutants through the Valley until reaching the mountains (Tehachapi Range) at the southern end of the basin, causing an eddy of air to form in the vicinity of Visalia and Fresno, just west of the southern Sierra Nevada. Frequent inversions trap the Valley air at night. Rising daytime air currents subsequently carry these trapped pollutants into the Sierra Nevada, resulting in some of the worst air quality found in any NPS unit in the country (Cahill et al.1996). This movement of polluted air into the Sierra occurs daily during summer months. The airflow pattern transports air pollutants to all the SIEN parks; the highest pollutant levels occur in Sequoia and Kings Canyon—lower, but not insignificant, levels occur in Yosemite and Devils Postpile.

Automobiles are one of the major sources of pollutants in the San Joaquin Valley, contributing much of the particulates, carbon monoxide, nitrogen oxides, and hydrocarbons annually emitted. The latter two constituents are precursors in the formation of ozone. Other sources of pollution include agricultural activities (e.g. plowing, diesel generators, off-road vehicles), power generation, petroleum production, and other industrial emissions.

Emissions within the parks include motorized vehicles, whose numbers are greatest in the summer when visitation is at its highest. Smoke emissions are another source found within park boundaries. Fires from prescribed burns, wildfires, and campfires all produce varying levels of particulate matter and chemical precursors to ozone. During winter months, most in-park emissions are from wood used for heating; however, emissions are typically minimal due to low

visitation and a smaller number of in-park residents. Still, a vast majority of air pollutants found in the parks are generated outside park boundaries.

There are also air-related health concerns in SIEN parks. Increased levels of ozone and particulate matter pose a threat to human health. The U.S. Environmental Protection Agency (USEPA) has established National Ambient Air Quality Standards (NAAQS) for each of six “criteria” pollutants, to protect the public from the health hazards associated with air pollution. These six criteria pollutants are: carbon monoxide (CO), ozone (O₃), nitrogen dioxide (NO₂), sulfur dioxide (SO₂), particulate matter less than 10 micrometers in diameter (PM₁₀), particulate matter less than 2.5 micrometers, and lead (Pb)(Table 1). The state of California has adopted additional air quality standards for sulfates, hydrogen sulfide, vinyl chloride, and visibility-reducing particulates. Ozone may pose a risk to animal health as well.

TABLE 1. National and California Ambient Air Quality Standards

Pollutant	Averaging Time	Federal Primary Standards	California Standards
Particulate Matter (PM10)	Annual Arithmetic Mean	50 ug/m3	20 ug/m3
	Annual Geometric Mean		30ug/m3
	24-hour	150 ug/m3	50 ug/m3
	1-hour	---	
Particulate Matter (PM 2.5)	Annual Arithmetic Mean	15 ug/m3	12 ug/m3
	Annual Geometric Mean	---	---
	24-hour	65 ug/m3	---
	1-hour		
Sulfur Dioxide (SO2)	Annual	80 ug/m3 (0.03ppm)	---
	24-hour	365 ug/m3 (0.14 ppm)	105 ug/m3 (0.04 ppm)
	3-hour		655 ug/m3 (0.25 ppm)
	1-hour	---	
Nitrogen Dioxide (NO2)	Annual	100 ug/m3 (0.053 ppm)	---
	1-hour	---	470 ug/m3 (0.25 ppm)
Ozone (O3)	8-hour	0.08 ppm	---
	1-hour	0.12 ppm (235 ug/m3)	0.09 ppm (180 ug/m3)
Carbon Monoxide (CO)	8-hour	9 ppm (10 mg/m3)	9.0 ppm (10 ug/m3)
	1-hour	35 ppm (40 mg/m3)	20 ppm (23 ug/m3)
Sulfates	24-hour	---	25 ug/m3
Lead	Calendar Quarter	1.5 ug/m3	---
	30-day average	---	1.5 ug/m3
Hydrogen Sulfide	1-hour	---	0.03 ppm (42 ug/m3)
Vinyl Chloride (chloroethene)	24-hour	---	0.010 ppm (26 ug/m3)

ug/m3 = micrograms per cubic meter

ppm = parts per million

--- = no standard exists for that pollutant and/or averaging time

Criteria Pollutants: Description and Issues

Air pollutants which are determined to have a detrimental effect on human health are called criteria pollutants.

Carbon Monoxide

Carbon monoxide (CO) is a colorless, odorless and poisonous gas produced by incomplete burning of carbon in fuels. When CO enters the bloodstream, it reduces the delivery of oxygen to the body's organs and tissues. Health threats are most serious for those who suffer from cardiovascular disease, particularly those with angina or peripheral vascular disease. Exposure to elevated CO levels can cause impairment of visual perception, manual dexterity, learning ability, and performance of complex tasks. Seventy-seven percent of the nationwide CO emissions are from transportation sources. The largest emissions contribution comes from highway motor vehicles. Thus, focus of CO monitoring has been on traffic-oriented sites in urban areas where the main source of CO is motor vehicle exhaust. Other major CO sources are wood-burning stoves, incinerators and industrial sources. The National Ambient Air Quality Standard (NAAQS) for carbon monoxide is a 9 ppm, 8-hour non-overlapping average, not to be exceeded more than once per year. An area meets the carbon monoxide NAAQS if no more than one 8-hour value per year exceeds the 9ppm threshold. To be in attainment, an area must: (1) meet the NAAQS for two consecutive years, and (2) carry out air quality monitoring during this entire time period. Air quality carbon monoxide value is estimated using EPA guidance for calculating design values ([Laxton Memorandum, June 18, 1990](#)).

Nitrogen Dioxide

Nitrogen dioxide (NO₂) is a brown-colored, highly reactive gas that is present in all urban atmospheres. NO₂ can irritate lungs, cause bronchitis and pneumonia, and lower resistance to respiratory infections. Nitrogen oxides are an important precursor both to ozone (O₃) and acid rain, and may affect both terrestrial and aquatic ecosystems. The major mechanism for the formation of NO₂ in the

atmosphere is the oxidation of the primary air pollutant nitric oxide (NO). Oxides of nitrogen (or NO_x) play a major role—together with volatile organic compounds (VOCs)—in atmospheric reactions that produce O₃ and PM 2.5 (PM=particulate matter). NO_x forms when fuel is burned at high temperatures. The two major emissions sources are transportation and stationary fuel-combustion sources, such as electric utility and industry boilers.

Ozone

Ozone (O₃) is a photochemical oxidant and the major component of smog. While O₃ in upper atmosphere is beneficial to life by shielding earth from harmful ultraviolet radiation from the sun, high concentrations of O₃ at ground level are a major health and environmental concern. O₃ is not emitted directly into the air, but is formed through complex chemical reactions between precursor emissions of volatile organic compounds (VOC) and oxides of nitrogen (NO_x) in the presence of sunlight. These reactions are stimulated by sunlight and temperature, so peak O₃ levels occur typically during the warmer times of the year. Both VOCs and NO_x are emitted by transportation and industrial sources. VOCs are emitted from sources as diverse as autos, chemical manufacturing, dry cleaners, paint shops and other sources that use solvents. The reactivity of O₃ causes health problems because it damages lung tissue, reduces lung function, and sensitizes the lungs to other irritants. Scientific evidence indicates that ambient levels of O₃ not only affect people with impaired respiratory systems, such as asthmatics, but healthy adults and children as well. Exposure to O₃ for several hours at relatively low concentrations has been found to significantly reduce lung function and induce respiratory inflammation in normal, healthy people during exercise. This decrease in lung function is generally accompanied by symptoms including chest pain, coughing, sneezing, and pulmonary congestion.

Particulate Matter

Air pollutants called particulate matter include dust, dirt, soot, smoke, and liquid droplets directly emitted into the air by sources such as factories, power plants, cars, construction activity, fires, and natural windblown dust. Particles formed in the atmosphere by condensation or the transformation of emitted gases such as sulfur dioxide (SO₂) and VOCs are also considered particulate matter. Another byproduct of atmospheric reactions is nitric acid (HNO₃), which combines with ammonia (NH₃) to produce a species of particulate matter (PM 2.5) called ammonium nitrate (NH₄NO₃). Based on studies of human populations exposed to high concentrations of particles (sometimes in the presence of SO₂) and laboratory studies of animals and humans, there are major effects of concern for human health. These include effects on breathing and respiratory symptoms, aggravation of existing respiratory and cardiovascular disease, alterations in the body's defense systems against foreign materials, damage to lung tissue, carcinogenesis, and premature death. The major subgroups of the population that appear to be most sensitive to the effects of particulate matter include individuals with chronic obstructive pulmonary or cardiovascular disease or influenza, asthmatics, the elderly, and children. Particulate matter also soils and damages materials, and is a major cause of visibility impairment in the United States. Annual and 24-hour National Ambient Air Quality Standards (NAAQS) for particulate matter were first set in 1971. Total suspended particulate (TSP) was the first indicator used to represent suspended particles in the ambient air. Since July of 1987, however, EPA has used—as an indicator—those particles with aerodynamic diameter smaller than 10 micrometers. Most recently, studies have found that even smaller diameter particles (under 2.5 micrometers or PM_{2.5}) are an even better indicator of health effects. These smaller particles are likely responsible for most adverse health effects from particulate matter because of their ability to infiltrate the thoracic or lower regions of the respiratory tract.

Sulfur Dioxide

High concentrations of sulfur dioxide (SO₂) affect breathing and may aggravate existing respiratory and cardiovascular disease. Sensitive populations include asthmatics, individuals with bronchitis or emphysema, children, and the elderly. SO₂ is also a primary contributor to acid deposition, or acid rain, which causes acidification of lakes and streams and can damage trees, crops, historic buildings, statues, etcetera. In addition, sulfur compounds in the air contribute to visibility impairment in large parts of the country. This is especially noticeable in national parks. Ambient SO₂ results largely from stationary sources such as coal and oil combustion, steel mills, refineries, pulp and paper mills and from nonferrous smelters. There are three NAAQS for SO₂: an annual arithmetic mean of 0.03 ppm (80 ug/m³); a 24-hour level of 0.14 ppm (365 ug/m³); and a 3-hour level of 0.50 ppm (1300 ug/m³). The first two standards are primary (health-related) standards, while the 3-hour NAAQS is a secondary (welfare-related) standard. The annual mean standard is not to be exceeded, while the short-term standards are not to be exceeded more than once per year.

Attainment

To reiterate, air pollutants determined to have a detrimental effect on human health are called criteria pollutants. Geographic areas are determined, by both national and California standards, as meeting or not meeting the health based standard for a specific air pollutant (e.g., ozone, PM₁₀). An area may be in attainment for one pollutant and non-attainment area for others. The following tables describe attainment status for Sierra Nevada network parks (Table 2). Non-attainment status requires that the area produce a plan, with a targeted timeline, to reduce pollutant levels within the specified area and submit the plan to the appropriate level of government (state or federal) for approval.

Table 2. Attainment status for criteria air pollutants in SIEN parks: Yosemite (YOSE)–Mountain Counties (Mariposa and Tuolumne) Air District; Sequoia & Kings Canyon National Parks (SEKI) and Devils Postpile National Monument (DEPO)–San Joaquin Valley Air District

Yosemite:

Pollutant	Federal	State
Yosemite		
O3 (1hr)	Attainment	Non-Attainment
O3 (8hr)	Non-Attainment	_____
CO	Attainment	Unclassifiable
PM10	Attainment	Non-Attainment
PM2.5	_____	Unclassifiable
SO2	Attainment	Attainment
NO2	Attainment	Attainment
Sequoia & Kings Canyon and Devils Postpile		
O3 (1hr)	Non-Attainment/Extreme	Non-Attainment
O3 (8hr)	Non-Attainment/Serious	_____
CO	Attainment	Attainment
PM10	Non-Attainment/Serious	Non-Attainment
PM2.5	_____	Non-Attainment
SO2	Attainment	Attainment
NO2	Attainment	Attainment

Table 3. Target dates for attainment (as of 17 May 2004)..

Air District	Pollutant	Averaging Time	Classification	Attainment Year
San Joaquin Valley	Ozone	1-hour	Extreme	2010
	Ozone	8-hour	Serious	2013
	Particulate Matter (PM10)		Serious	***
Mariposa/Tuolumne	Ozone	1-hour	***	***
	Ozone	8-hour	Subpart 1	2009
	Particulate Matter (PM10)			

*** no data available and/or no attainment classification

Air Quality in Sierra Nevada Parks

Emissions:

Devils Postpile and Sequoia and Kings Canyon are located within the San Joaquin Valley Unified Air Pollution Control District (SJVUAPCD). The SJVUAPCD is made up of 8 counties (San Joaquin, Stanislaus, Merced, Madera, Fresno, Tulare, Kings, and Kern) in the San Joaquin Valley. This air district is in non-attainment status for ozone and PM10 (Table 2).

Yosemite is within Mountain Counties' air district, comprising Tuolumne and Mariposa Counties. A small portion of southern YOSE is located in Madera County but the park is generally not regulated by the SJVUAPCD. Mountain Counties Air District also is in non-attainment status for ozone and PM10 (Table 2).

All SIEN parks are exposed to pollutants transported from the Valley as well as from other urban areas of California due to prevailing westerly winds. Approximately nine percent of the state's thirty five million people live in San Joaquin Valley (SJV), and emission sources (e.g. industry, agriculture, and transportation) within SJV account for about 14 percent of total statewide emissions (Alexis et al. 1999).

Emissions in SJV derive from a number of moderate-sized urban areas, primarily located along the Highway 99 corridor. The other main travel corridor in SJV is US Interstate 5. Since 1980, population growth in the SJV has been more rapid than in other parts of California, partially offsetting effects of emission-control programs (Alexis et al. 1999). Air pollution sources found within parks include: automobiles, buses, construction activities, campfires, wood-burning stoves, natural and prescribed fire, and road dust.

Principal air pollutant species of concern to SIEN parks are ozone, ozone precursors (NOX and VOC's), particulate matter precursors, and particulate matter concentrations that decrease visibility and/or lead to acidic deposition. SO₂ emissions are not high in any SIEN parks. Air districts' target dates for ozone and PM10 standards attainment are shown in Table 3.

Sequoia and Kings Canyon are located within Fresno and Tulare counties. No major point sources of emissions are located in Tulare County; major point sources are not numerous in Fresno County, nor in most other counties of the SJV. Within Fresno County, the closest sources that emit at least 100 tons/year of VOC, NO_x, PM10, or SO₂ are located near Fresno and Kingsburg; three additional point sources are located in the western portion of the county. As of 1996, stationary sources accounted for 22 % of ROG emissions, 34 % of NO_x emissions, and 4% of PM10 emissions in Fresno County (CARB, 1998b). In Tulare County, stationary sources accounted for 8 % of ROG emissions, 11 % of NO_x emissions, and 5 % of PM10 emissions in 1996 (CARB, 1998b). In both Fresno and Tulare counties, mobile sources dominated NO_x emissions, while area sources (road dust, construction, and farming operations) dominated PM emissions. Mobile sources accounted for 46 % of VOC emissions in Fresno and 41 % in Tulare Counties.


Yosemite is located within Madera, Mariposa, and Tuolumne counties. Major point sources are not numerous in these counties, nor in most other counties of the SJV or Mountain Counties air basins. Sources that emit at least 100 tons/yr of VOC, NO_x, PM10, or SO₂ are located near Jamestown and

Standard in Tuolumne County, and further away, near Madera and Chowchilla in western Madera County; none are located within Mariposa County. As of 1996, stationary sources accounted for 12% of VOC emissions, 36% of NOx emissions, and 7% of PM10 emissions in Madera, Mariposa, and Tuolumne counties (CARB1998b). Mobile sources dominated NOx emissions, whereas area sources (road dust, construction, and farming operations) dominated PM emissions.

Devils Postpile is technically in SJVUAPCD, but is located at the extreme eastern edge of Madera County. It straddles the San Joaquin Valley (SVJ) and Great Basin (GB) Air Basins. There are significant differences between the two air basins. One is size: the SVJ is larger, comprising 8 counties; the GB comprises 3 counties. There are both ozone and particulate matter concerns in SVJ; in GB, particulate matter is more prominent. Area wide sources are the main contributor of PM for both districts. Mobile sources are the most influential in contributing precursors of ozone in SVJ. Monitoring near the monument provides air-related data that are relevant to DEPO.

Ozone:

Through monitoring and research at Sequoia, Kings Canyon and Yosemite, we know that ozone levels are consistently higher than background or natural levels. All three parks experience multiple days where ozone levels exceed the state and national health standard (for 8hr ozone). While high levels of ozone can be found at all of the Class I SIEN parks, levels are consistently and at times substantially higher at Sequoia and Kings Canyon. For example, in 2003 Sequoia and Kings Canyon exceeded the health standard for ozone (85 ppb 8 hr. average) 72 days, while Yosemite recorded 10 days above the standard. These numbers of days exceeding health standards ranks Sequoia, Kings Canyon and Yosemite in the top 10 most ozone polluted parks in the country.



Ozone is a pollutant well known for producing foliar damage in ozone sensitive ponderosa and Jeffrey pines. Ozone damages photosynthetic cells in these pines. It impairs their ability to grow, and compete with other plants, and compromises their resistance to other forest stressors such as insects, pathogens, and drought. (James 1980, Miller 1983). Monitoring sensitive vegetation for ozone damage will help define the severity of impact to our resources. Potential bioindicator plant species, plants with sensitivity to ozone pollution for the SIEN parks are listed in Table 4 (NPS - Air Resources Division).

Table 4. Bioindicator plant species

Park	Scientific Name	Common Name
SEKI,YOSE	Apocynum androsaemifolium	Spreading dogbane
SEKI,YOSE	Artemisia douglasiana	Mugwort
SEKI,YOSE	Physocarpus capitatus	Ninebark
SEKI,YOSE	Pinus jeffreyi	Jeffrey pine
SEKI,YOSE	Pinus ponderosa	Ponderosa pine
SEKI,YOSE	Populus tremuloides	Quaking aspen



SEKI,YOSE	<i>Rhus trilobata</i>	Skunkbush
SEKI,YOSE	<i>Salix scouleriana</i>	Scouler's willow
SEKI,YOSE	<i>Sambucus mexicana</i>	Blue elderberry

Nitrogen and Sulfur Deposition:

Although acidic deposition has been found to currently have less significant impact on Sierra Nevada resources, there are two seasonal pulses of acidity: (1) late summer rain, and (2) the onset of spring snowmelt. These pulses of relatively high acidity can lower pH below 5.35, making naturally basic waters acidic in an un-buffered ecosystem (i.e. not much vegetative cover, steep slopes, granite under-layer, not much soil) (Melack and Sickman et al. 1998) Current information shows that these high elevation systems are more resilient than originally believed (Melack and Sickman et al. 1998). Currently, these systems are able to buffer existing acidic events. Potential reactions of these systems to increased acidic deposition are not well understood. In forested ecosystems, summer wet deposition concentrations of NO₃⁻ and SO₄⁻ averaged two- and five-times higher, respectively, as compared with concentrations reported for remote areas of the world (Stohlgren and Parsons 1987). Though it is uncertain what minimum concentrations of acid may cause direct or indirect foliar damage to plant species in Sierra Nevada parks, Stohlgren and Parsons (1987) emphasized that it is important to note that acid concentrations are highest during the primary growing season and the period of maximum drought stress, both of which occur during the summer (Stohlgren and Parsons 1987; Williams and Melack 1991; Williams and Melack 1997b; Melack et al. 1998; Williams et al. 1995; Galloway et al. 1982; Stoddard 1994)

Wet deposition monitoring has provided a long term data set in Sequoia, Kings Canyon, and Yosemite, measuring constituents that are related to vehicle, agriculture, and other human activities which produce air pollutants. The data show that Parks in the West do not receive substantial amounts of wet SO₄ deposition. On the other hand, levels of NO₃ and NH₄ deposition appear high

enough to fertilize terrestrial vegetation and cause eutrophication of aquatic systems. Without site specific dry deposition data however, a quantitative (i.e., determination of critical loads) approach is impossible. Figure 1 illustrates approximately the past 20 years of annual deposition of NO_3 , NH_4 , and SO_4 for Sequoia, Kings Canyon, and Yosemite. These numbers do not provide a complete picture of N contribution from dry deposition in the parks: given their Mediterranean climate, almost half the year is dry. Therefore wet deposition values alone underestimate total deposition.







Figure 1. Total Annual Wet Deposition (NADP), SEKI and YOSE, 1982 – 2003.

Particulate Matter/Visibility:

Changes in visibility at SIEN parks is largely due to organics, sulfates, and nitrates. Historically, visibility varies with patterns in weather, winds (and the effects of winds on coarse particles), and smoke from fires. No information is available on how distribution of present visibility conditions differs from the profile under "natural" conditions, but the cleanest 20% of days probably approaches natural conditions (GCVTC 1996). Smoke from frequent fires is suspected to have reduced pre-settlement visibility below current levels during some summer months. Long-term visibility trends fall into three categories: increases, decreases, and insignificant changes. The characterization of long-term trends can be a highly subjective exercise: slopes and their statistical significance can vary depending on the technique employed. Recently the IMPROVE aerosol network, initiated in March 1988, matured to a point where long-term trends of

average ambient aerosol concentrations and reconstructed extinction can be assessed for sites with an eleven year period of record. (CA Class I Parks 2001).

Because full IMPROVE aerosol monitoring was not initiated at Yosemite or Sequoia until March 1994, the current monitoring period is not considered long enough to adequately summarize long-term trends. Seasonal trends (1996 to 1998) show that fine aerosols rise in the summer and that organics make up the largest proportion of these aerosols (Fig.2; <http://vista.circa.colostate.edu/improve/Data/GraphicViewer/>). other aerosol species observed include: sulfate, nitrate, elemental (light absorbing) carbon, and coarse particles (i.e., wind blown dust). Coarse particles tend to have natural origins and deposit close to their source. Fine particles are usually man-made, have the ability to transport great distances, and cause the greatest impairment to visibility.






Figure 2. Fine aerosol mass proportions for Yosemite and Sequoia & Kings Canyon National Parks, 1996-1998. Light absorbing carbon noted as LAC, above.

In the Malm et al. (2000) IMPROVE report, eleven years of data were used to describe trends in particulate matter (specifically, PM 2.5) pollution for Yosemite (CA Class I parks 2001). Aerosol species were associated with low, median, and high fine mass day categories, and then respective light extinctions and deciviews (relative haziness values) were calculated for each species. The report concluded that there was an insignificant change in visibility over time, meaning that there was neither an increasing nor a decreasing trend (Fig.3). However, like Sequoia, seasonal trends within Yosemite show a significant rise in fine aerosols through the hot summer months, with organics making up the largest proportion of these aerosols. Visibility diminishes with increased particulate matter pollution, so the IMPROVE network attempts to identify chemical species that significantly reduce visibility in our Parks.

By summarizing these data regionally, and comparing them nationally, it becomes apparent that southern California visibility is similar to northern Massachusetts and southern Vermont in terms of average haziness in deciviews (Fig. 3). However, approximately 11% of southern California's light absorption is caused by aerosols, while only 5.1% of New England's light absorption is caused by aerosols. This is a complicated observation considering the differences in

average air moisture content between the two regions, as well as the variety of pollution species, all contributors to haze. The worst average visibility in the country is in the southeast, with visibility impairment values of 21 to 26 deciviews (lower numbers denote greater visibility).




Figure 3: Regional visibility measured in deciviews, 1996 to 1998. Lower numbers denote greater visibility.

Contaminants

California pesticide-use data show that between 1991 and 2000, approximately 2 billion pounds of active ingredients were applied in the State (Pesticide Action Network). Pesticides, which are used extensively in California's heavily-farmed SJV, are of concern in the Sierra Nevada. Potential impacts depend on the degree to which a particular pesticide breaks down in the environment and the levels of toxicity of the breakdown products. The amount of

pesticide use suggests that further study is warranted. Pesticides vary as to their levels of volatility. For example, in Fresno County, the estimated pesticide VOC emissions potential of 17.6 million pounds (8.8 thousand tons) is a substantial portion of the total VOC emissions of 39.4 thousand tons. However, California Department of Pesticide Regulation's (CDPR) estimate of VOC emissions potential exceeds that used in the CARB emissions inventory (CARB, 1998b) by a factor of two.

Pesticides are of concern for their potential toxicity, not just their ozone-forming potential. Studies conducted in SEKI indicate that more commonly and heavily used pesticides in the SJV are transported into the Sierra Nevada (Zabik and Seiber 1993, LeNoir et al. 1999). Measurements made in Sequoia National Park, showed pesticides concentrations diminishing between Ash Mountain entrance station and Lower Kaweah (Zabik and Seiber 1993). This represents a decrease in concentration with both distance and elevation (relative to the San Joaquin valley floor). Pesticides applied to crops (that are intensively produced in the Central Valley) can volatilize under warm valley temperatures, subsequently be transported via upslope air movement, and then deposited in the cooler, high-elevation regions of Sequoia National Park (Zabik and Seiber 1993; LeNoir et al. 1999).

In 2002, SEKI was included in the Mercury Deposition Network (MDN), a network developed by the National Atmospheric Deposition Program (NADP). The sampler is co-located with the NADP sampler at Lower Kaweah in Sequoia National Park. This network investigates the local and regional transport and deposition of mercury. Mercury is known to contaminate aquatic ecosystems, and there are over 30 states which have fish consumption advisories due to mercury content. Atmospheric transport and deposition is the dominant pathway to aquatic ecosystems. This problem has greater attention in the eastern U.S., as reflected in both the number of sampling sites in the east, and mercury levels found in these states. Origin of mercury in the East is predominantly from coal burning industries. There also exists global transport of mercury in the atmosphere, whose source is wide and varied. Effects of mercury deposition vary

by site as a result of differences in soil chemistry, water chemistry, the food web, etc. There is less information on mercury deposition in California, and there are only 2 active MDN sites in California.

Air Resources Division of the National Park Service

The National Park Service Air Resources Division (ARD) mission is to: “[p]reserve, protect, enhance, and understand air quality and other resources sensitive to air quality in the National Park System” (<http://www2.nature.nps.gov/air/>). In order to do this, a combination of both monitoring and research is necessary. Long-term monitoring data make it possible to plot and assess air pollution patterns and compare these patterns to other regions of the country. At the same time, research data are needed to answer more specific pollution questions concerning cause and effect and total deposition loads.

As part of the NPS ozone monitoring network, and by participating in the National Atmospheric Deposition Program (NADP), the Interagency Monitoring of Protected Visual Environments (IMPROVE), and the Clean Air Status & Trends Network (CASTNet), Sequoia & Kings Canyon and Yosemite National Parks contribute to the NPS effort to monitor air pollution in the form of wet deposition and dry deposition, as well as fine particulate matter and aerosols. All of these monitoring data are collected with relevant meteorological data, making the interpretation of the pollutant patterns more meaningful and comparable. Air related research projects are mostly cooperative with the NPS in the role of facilitator, supplying information and resources to universities, private researchers, and other branches of state and federal government.

Long-term Monitoring Programs:

The National Park Service Air Resource Division (ARD) coordinates NPS air quality monitoring programs nationwide and contracts with a private consulting group called Air Resources Specialists (ARS) located in Fort Collins, Colorado. Specifically, ARS provides technical support which enables NPS

station operators to troubleshoot complicated monitoring equipment and provide for consistent operations. Their efforts contribute to generation of accurate and high quality data. The ARS group also helps NPS-ARD with data summaries and interpretation.

Both Yosemite and Sequoia & Kings Canyon operate year-round air monitoring sites, some dating back to the early 1980s (Fig. 4). Each site is unique in its array of monitoring equipment (described in Table 5). Devils Postpile National Monument does not currently conduct any air monitoring within the boundaries of the monument, but monitoring work has been conducted in the surrounding area and is representative of air in the monument.





Figure 4. Air quality monitoring sites in Sequoia & Kings Canyon and Yosemite National Parks.

Table 5. Air monitoring stations

Park	Site	Type	Network **
Sequoia	Ash Mountain	Meteorology	NPS – ARD
		Ozone	NPS – ARD
		Particulate Matter	SEKI Fire
		PM 2.5 and 10	IMPROVE
	Lookout Point	Meteorology	NPS – ARD
		Ozone	NPS – ARD
		Dry deposition	CASTNet
	Lower Kaweah	Meteorology	NPS – ARD
		Ozone	NPS – ARD
		Wet deposition	NADP/NTN
		Mercury	MDN
		Webcam	NPS – ARD
Yosemite	Yosemite Valley	PM 2.5 and 10	CARB
		Meteorology	NPS – ARD
		Ozone	NPS – ARD
		NOX	NPS – ARD
		CO	NPS - ARD
	Turtleback Dome	Meteorology	NPS – ARD
		Ozone	NPS – ARD
		Visibility	NPS – ARD
		Webcam	YOSE Concession
		Dry deposition	CASTNet
		Particulate Matter	IMPROVE
	Hodgdon Meadow	Wet deposition	NADP/NTN

- ** California Air Resources Board (CARB)
- Clean Air Status and Trends Network (CASTNet)
- Interagency Monitoring of Protected Visual Environments (IMPROVE)
- Mercury Deposition Network (MDN)
- National Atmospheric Deposition Program/National Trends Network (NADP/NTN)
- National Park Service – Air Resources Division (NPS-ARD)

National Atmospheric Deposition Program/National Trends Network

National Atmospheric Deposition Program/National Trends Network

(NADP/NTN) is a nationwide network of over 200 precipitation monitoring sites developed by the EPA. The network is a cooperative effort between many different groups, including the State Agricultural Experiment Stations, U.S. Geological Survey, U.S. Department of Agriculture, and numerous other governmental and private entities. The purpose of the network is to collect weekly rain, snow, or condensation samples and produce data on its chemistry. The weekly water samples are sent to the Central Analytical Laboratory where they are analyzed for hydrogen (acidity as pH), sulfate, nitrate, ammonium, chloride, and base cations (such as calcium, magnesium, potassium and sodium (*paraphrased from <http://nadp.sws.uiuc.edu/nadpoverview.asp>*).

The NADP site within Giant Forest, Sequoia National Park, has been in operation since the early 1980's, while the NADP site at Hodgden Meadows in Yosemite has been operating since 1981.

Interagency Monitoring of Protected Visual Environments

Interagency Monitoring of Protected Visual Environments (IMPROVE) is a cooperative measurement effort comprising representatives from federal and regional-state organizations. It aids the creation of federal and state implementation plans for the protection of visibility in Class I areas (156 national parks and wilderness areas) as stipulated in the 1977 amendments to the Clean Air Act .The objectives of IMPROVE are:

- (1) to establish current visibility and aerosol conditions in mandatory class I areas;
- (2) to identify chemical species and emission sources responsible for existing man-made visibility impairment;
- (3) to document long-term trends for assessing progress towards the national visibility goal; and
- (4) and with the enactment of the Regional Haze Rule, to provide regional haze monitoring representing all visibility-protected federal class I areas where practical.

IMPROVE has also been a key participant in visibility-related research, including the advancement of monitoring instrumentation, analysis techniques, visibility modeling, policy formulation and source attribution field studies (*paraphrased from <http://vista.circa.colostate.edu/improve/Overview/Overview.htm>*).

A full IMPROVE aerosol sampler (modules A, B, C, and D) began operation at Sequoia in March of 1994. The sampler is located near Park Headquarters at Ash Mountain, at the southwest entrance to Sequoia National Park. There has been a full IMPROVE aerosol sampler at Yosemite since March 1988, located at Turtleback Dome. In 2002 improvements to the Sequoia IMPROVE stations added a set of PM10 modules (E & F) at the Ash Mountain site.

Clean Air Status and Trends Network

Clean Air Status and Trends Network (CASTNET) was developed by EPA to monitor dry acid deposition. By design, monitoring site locations are predominantly rural to assess the relationship between regional pollution and changes in regional patterns in deposition. CASTNET also includes measurements of rural ozone and the chemical constituents of PM 2.5 (fine particulate matter). Rural monitoring sites of CASTNET provide data where sensitive ecosystems are located and provide insight into natural background levels of pollutants where urban influences are minimal. These data provide

much-needed information to scientists and policy analysts to study and evaluate numerous environmental effects, particularly those caused by regional sources of emissions for which long range transport plays an important role. Measurements from these networks are also important for understanding non-ecological impacts of air pollution such as visibility impairment and damage to materials, particularly those of cultural and historical importance (*paraphrased from <http://www.epa.gov/castnet>*).

The CASTNet dry deposition monitoring site located at Lookout Point, within Sequoia National Park, began operating February 1997. The CASTNet site in Yosemite, on Turtleback dome, has been in operation since September 1995.

Light Pollution

Light pollution is not confined to cities. Excessive glare, urban sky glow, and poor lighting threaten dark skies. Night sky visibility is an important aesthetic value of wilderness and its protection has been added to the responsibilities of NPS managers. The objectives of the night visibility monitoring program are; 1) define the magnitude of night sky brightness, 2) identify short-term or seasonal impairments, 3) utilize monitoring data summaries for public education, and 4) help define limits of acceptable change for night sky visibility.

Dark sky monitoring sites were established in both Sequoia and Kings Canyon. Yosemite currently has one site with a second identified and monitoring began in September 2004.

Natural Soundscape

Biophony is the combined sound that living organisms produce in a given habitat. When habitat health is stable, indications are that there is a symbiotic and unique relationship between the creature voices denoted by both frequency and time niches discernible on audio spectrogram graphic displays with a high degree of discrimination and detail (Krause 2001). Manmade noise can be harmfully disruptive and, as such, NPS has designated natural soundscapes as a resource, which means they must be protected (akin to “clean air”). The program is managed by the NPS-ARD and parks locally facilitate and conduct sound measurements.

In 2001 Sequoia National Park became the site of a natural soundscape monitoring pilot program. Dr. Bernie Krause and Dr. Stuart Gage collected recordings in different habitats at different seasons to test biophony as an indicator of habitat fitness.

Monitoring and Research Needs

Data Acquisition and Analysis, Monitoring

Current monitoring efforts need to continue. In addition to monitoring efforts, resources should also be devoted to acquiring and integrating existing data, and to analyze and present findings from the available datasets. Existing meteorological and air-quality data could be more extensively analyzed to better characterize patterns of transport of air pollutants (e.g., fine particulate), including the nature of upslope and downslope flows.

Better characterization of pollutant transport is also needed to support the parks' fire management programs, help discern origins of specific pollutants, and provide critical information to complement biological effects research.

Monitoring of particulate matter needs to be updated, modified, and coordinated with adjacent land managers. Emissions from wildland fires are not the largest source of particulate matter, but the smoke produced can temporarily degrade air quality on a local and/or regional scale. Expansion of the PM monitoring program is necessary to determine extent of smoke transport and effects to human health. Smoke produces particulate matter (PM₁₀ and PM_{2.5}), CO, and precursors to ozone, potentially affecting park employees, visitors, and residents of nearby communities. Additional PM monitors, both stationary and mobile are needed to adequately assess levels for park burning operations and provide real-time data for health advisories. During large smoke events carbon monoxide (CO) monitoring should be considered. Monitoring locations should include sites near campgrounds, offices, residences and in neighboring communities. Real-time data displays (e.g., daily particulate levels) are potentially of significant interest to park visitors.

Additional efforts should be devoted to characterizing the ambient atmospheric levels of non-criteria air pollutants (e.g., pesticides) and comparing these levels with their concentrations in aquatic systems. This research effort would be enhanced by linking it to any ongoing studies of atmospheric transport and biological effects.

Wet and Dry Deposition

Continued monitoring of wet and dry deposition is needed to provide estimates of spatial patterns and temporal variability of deposition. The long-term data records should be maintained by continuing wet-deposition monitoring at Giant Forest (NADP). Dry deposition monitoring should continue at Ash Mountain (IMPROVE), and Lookout Point (CASTNet). It is recommended that wet deposition monitoring be reinitiated at Ash Mountain in Sequoia National Park.

The long-term deposition record presently available lends itself to analyses of trends. The data should be analyzed to determine the statistical power of trend tests (i.e., the length of monitoring needed to detect trends and the levels of change that can be discerned within a specified time).

Increased input of nitrogen (i.e. HNO_3 , NH_4 , NH_3 , NO_3) is recognized as an additional stressor to Sierra Nevada ecosystems. There are potential landscape scale changes such as terrestrial fertilization and aquatic eutrophication due to increased nitrogen and phosphorus loading (Sickman et al. and Fenn et al.) Improved measurements of both wet and dry forms of nitrogen will give more accurate levels of nutrient loading to Sierra Nevada ecosystems.

Additional benefit to understanding pollutant levels in the Sierra Nevada would be use of NO_x analyzers. Co-location with ozone, wet deposition, and particulate matter would be useful to park management needs, including but not limited to fire operations, health advisories, and monitoring vegetation change.

Ozone

Continuous measurements of ozone are needed to provide estimates of spatial patterns and temporal variability. The long-term data records should be maintained by continuing ozone monitoring at the existing sites at Giant Forest, Ash Mountain, and Lookout Point in Sequoia and at Turtleback Dome in Yosemite. An ozone monitor was installed in Yosemite Valley to provide information to an ozone effects study in 2001. The study found significant differences in ozone levels between the Valley site and Turtleback Dome. With

the high visitation found in the Valley, this site should be retained and added to the NPS ozone monitoring network.

Passive ozone monitors suggest that ozone levels vary significantly within Sierra Nevada parks. Additional efforts to characterize the spatial patterns of ozone concentrations should be supported. Summary statistics documenting violations of the state and federal ozone standards are also of interest. The use of existing passive sampler data should be continued to provide on-site data relevant for co-located ozone-injury vegetation plots. The value of previous ozone data collected with passive samplers should be assessed for developing calibrated spatial relationships relative to analyzers, so that large-scale ozone exposure in the park can be quantified. If the spatial distribution (especially variation in elevation) has not been sufficient, it is recommended that a one-time study with well-distributed passive samplers be used for spatial calibration.

Particulate Matter/Visibility

IMPROVE aerosol monitoring should continue at Sequoia and Yosemite. As part of the IMPROVE Network expansion, the EPA and NPS have upgraded the aerosol sampler in Sequoia to include 2 additional modules for PM 10. While the site at Ash Mountain (550 m) serves the purpose of measuring what PM is traveling into the park via the Kaweah River drainage, it inadequately represents the average elevation of the parks. An additional site has been proposed, to be located at a higher elevation (2,250 m). This new location may better represent conditions for the high elevation landscape that dominates the Sierra Nevada. Aerosol concentrations can vary at different elevations throughout the parks. Aerosol and optical monitoring at both high and low elevations would provide information concerning the extent and nature of visibility impairment at Sierra Nevada parks. Emissions from industrial processes and wildland fires also contribute to visibility impairment in all SIEN parks. In addition to monitoring for human health concerns, monitoring should be developed to identify carbon emissions from industrial processes and wildland fires.

Visibility as measured by transmissometers is being utilized in Yosemite. The transmissometer receiver is located on Turtleback Dome with the target transmitter at Foresta. Additional visibility data should be considered in high visitation areas where broad vistas are a hallmark for their respective parks to measure degradation or improvement.

Natural Soundscapes

Natural sound monitoring should be developed for one or more of the SIEN parks. Co-location with a year round air monitoring site would be the most useful as sound levels could correspond with meteorology and various high air pollution events (e.g. high levels of particulate matter or ozone). It would also be logistically practical as the air monitoring sites are visited on a weekly basis.

Dark Sky

The NPS Dark Sky monitoring program is widespread and growing. Yosemite, Sequoia and Kings Canyon have areas targeted for repeat dark sky photography. Comparisons should be made between natural dark sky and light pollution often found in developed areas. This would help in guiding park operations in types and location of light fixtures and could provide information useful to lighting policy and planning for local communities.

References

- Alexis A., P. Gaffney, C. Garcia, C. Nystrom, and R. Rood. 1999. The 1999 California Almanac of Emissions and Air Quality. California Air Resources Board. Sacramento, CA. 310 pp.
- Cahill, T.A., J.J. Carroll, D. Campbell, and T.E. Gill 1996. Air quality. In: Status of the Sierra Nevada, Sierra Nevada Ecosystem Projects. Centers for Water and Wildland Resources, University of California, Davis. Pp. 1227-1261.
- CARB. 1998b. Emission Inventory 1996. California Air Resources Board. Sacramento, CA.
- Clean Air Status and Trends Network - <http://www.epa.gov/castnet>
- Cooperative Institute for Research in the Atmosphere—
<http://vista.circa.colostate.edu/improve/Overview/Overview.htm>,
<http://vista.circa.colostate.edu/improve/Data/GraphicViewer/>
- Fenn, M.E., R. Haeuger, G.S. Tonnesen, J.S. Baron, S. Grossman-Clark, D. Hope, D.A. Jaffe, S. Copeland, L. Geiser, H.M. Rueth, and J.O. Sickman 2003. Nitrogen emission, deposition, and monitoring in the Western United States. BioScience Vol. 53 No. 4
- Fenn, M.E., J.S. Baron, E.B. Allen, H.M. Rueth, K.R. Nydick, L. Geiser, W.D. Bowman, J.O. Sickman, T. Meixner, D.W. Johnson, and P. Neitlich 2003. Ecological effects of nitrogen deposition in the western United States. BioScience Vol. 53 No. 4
- Galloway, J.N., G.E. Likens, W.C. Keene, and J.M. Miller. 1982. The composition of precipitation in remote areas of the world. J. Geophys. Res. 88:10850-10864.
- Grand Canyon Visibility Transport Commission (GCVTC), 1996, Proposed Recommendations of the Grand Canyon Visibility Transport Commission, Public Advisory Committee Final Report.

- James R.L., F.W. Cobb, P.R. Miller, J.R. Parmeter Jr. 1980 Effects of oxidant air pollution on susceptibility of pine roots to *Fomes annosus*. *Phytopathology* 70:560-563.
- LeNoir, J.S., L.L. McConnell, G.M. Fellers, T.M. Cahill, and J.N. Seiber. 1999. Summertime transport of current-use pesticides from California's Central Valley to the Sierra Nevada Mountain Range, USA. *Environ. Toxicol. Chem.* 18(12):2715-2722.
- Miller, P.R. 1983. Ozone effects in the San Bernardino National Forest. In *Proceedings: Air Pollution and the Productivity of the Forest*. Izaak Walton League and Penn State University, pp. 161-197.
- Melack, J.M., J.O. Sickman, A. Leydecker, and D. Marrett. 1998. Comparative analyses of high-elevation lakes and catchments in the Sierra Nevada: susceptibility to acidification. Final Report, Contract A032-188, California Air Resources Board, Sacramento. 615 pp.
- National Atmospheric Deposition Program -
<http://nadp.sws.uiuc.edu/nadpoverview.asp>
- National Park Service—<http://www2.nature.nps.gov/air/Pubs/index.htm>
- Sequoia and Kings Canyon National Park—<http://www.nps.gov/seki/>
- Sickman, J.O., A. Leydecker, C.C.Y. Chan, C. Kendall, J.M. Melack, D.M. Lucero, J.P. Schimel 2003a. Mechanisms underlying export of nitrogen from high elevation catchments during seasonal transitions. *Biogeochemistry* 64:1-24
- Sickman, J.O, D. Clow, and J.M. Melack 2003b. Evidence for nutrient enrichment of high elevation lakes in the Sierra Nevada, California. *Limnology and Oceanography* 48(5) 1885-1892.
- Stoddard, J.L. 1994. Long-term changes in watershed retention of nitrogen: its causes and aquatic consequences. In: L.A. Baker, (ed.). *Environmental*

- Chemistry of Lakes and Reservoirs. Advances in Chemistry Series, No. 237, American Chemical Society, Washington, DC. pp. 223-284.
- Stohlgren, T.J. and D.J. Parsons. 1987. Variation of wet deposition chemistry in Sequoia National Park, California. *Atmos. Environ.* 21(6):1369-1374.
- Sullivan, T.J., Peterson, D.L., and C.L. Blanchard 2001. Assessment of Air Quality and Air Pollutant Impacts in Class I National Parks of California 377p.
- Williams, M.W. and J.M. Melack. 1997b. Atmospheric deposition, mass balances, and processes regulating streamwater solute concentrations in mixed-conifer catchments of the Sierra Nevada, California. *Biogeochemistry* 37:111-144.
- Williams, M.W., R.C. Bales, A.D. Brown, and J.M. Melack. 1995. Fluxes and transformations of nitrogen in a high-elevation catchment, Sierra Nevada. *Biogeochemistry* 28:1-31.
- Yosemite National Park—<http://www.nps.gov/yose/>
- Zabik, J.M. and J.N. Seiber. 1993. Atmospheric transport of organophosphate pesticides from California's Central Valley to the Sierra Nevada Mountains. *J. Environ. Qual.* 22:80-90.